MPP-2008-14,TIFR/TH/08-04

arXiv:0802.1481v3 [hep-ph] 20 Oct 2008

Identifying neutrino mass hierarchy at extremely small θ_{13} through Earth matter effects in a supernova signal

Basudeb Dasgupta,[1] Amol Dighe,[1] and Alessandro Mirizzi[2]

[1] *Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005, India*
[2] *Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Föhringer Ring 6, 80805 München, Germany*

Collective neutrino flavor transformations deep inside a supernova are sensitive to the neutrino mass hierarchy even at extremely small values of θ_{13}. Exploiting this effect, we show that comparison of the antineutrino signals from a galactic supernova in two megaton class water Cherenkov detectors, one of which is shadowed by the Earth, will enable us to distinguish between the hierarchies if $\sin^2\theta_{13} \lesssim 10^{-5}$, where long baseline neutrino experiments would be ineffectual.

PACS numbers: 14.60.Pq, 97.60.Bw

After many years of experiments on atmospheric, solar and terrestrial neutrinos, our knowledge of neutrino masses and mixing has grown by leaps and bounds [1, 2, 3, 4]. The data is now described satisfactorily in the 3-neutrino oscillation framework defined by two mass squared differences $\Delta m^2_\odot$ and $\Delta m^2_{\rm atm}$, three mixing angles θ_{12}, θ_{23} and θ_{13}, and the CP-violating phase δ. The parameters θ_{13} and $|\Delta m^2_{\rm atm}|$ are determined by atmospheric neutrino experiments and long baseline experiments, while $\Delta m^2_\odot$ and θ_{12} are determined by solar and reactor experiments. Current data on neutrino oscillations do not determine the sign of $\Delta m^2_{\rm atm}$. One refers to $\Delta m^2_{\rm atm} > 0$ as normal mass hierarchy and $\Delta m^2_{\rm atm} < 0$ as inverted mass hierarchy. For θ_{13} there is only an upper bound $\sin^2\theta_{13} < 0.02$ at 90 % C.L. from reactor experiments. The phase δ is completely unknown.

The primary goals of next generation neutrino experiments include the measurement of the leptonic mixing angle θ_{13}, and the determination of the neutrino mass hierarchy. The present and near-future long baseline and reactor experiments are expected to probe θ_{13} down to $\sin^2\theta_{13} \lesssim 10^{-2} - 10^{-3}$ (see, e.g., [5]). Over the next decade, if a neutrino factory or a superbeam facility is built, the reach could get extended to $\sin^2\theta_{13} \lesssim 10^{-4} - 10^{-5}$ [5, 6]. Several studies have been performed to ascertain the optimal strategy for determining the neutrino mass hierarchy if θ_{13} is large enough to be measured at the upcoming experiments (see, e.g., [7, 8, 9, 10]). However, if θ_{13} is small, then the determination of the mass hierarchy becomes very challenging even with next-to-next generation neutrino factory experiments [11, 12].

In this Letter, we propose a new astrophysical method for determining the neutrino mass hierarchy, which works for extremely small values of θ_{13} where the standard oscillation experiments fail. This method makes use of the Earth matter effects on the neutrino signal from a galactic supernova (SN).

Mikheyev-Smirnov-Wolfenstein (MSW) flavor conversions [13, 14] in a SN envelope are known to mix the primary neutrino fluxes. This mixing is crucially dependent on θ_{13} and the neutrino mass hierarchy. In addition to flavor conversions in the star, neutrinos arriving from a SN can undergo oscillations inside the Earth before being observed at a shadowed detector [15, 16, 17]. These Earth effects associated with supernova neutrinos have been studied extensively as a powerful tool to probe neutrino mass hierarchy for $\sin^2\theta_{13} \gtrsim 10^{-3}$ [18, 19, 20, 21]. For $\sin^2\theta_{13} \lesssim 10^{-5}$, the traditional analysis of neutrino flavor conversions in SN, where only MSW transitions were taken into account, predicts no hierarchy dependence of SN neutrino spectra arriving at the Earth [22]. Therefore, observation of the Earth matter effect cannot determine the neutrino mass hierarchy in that case. This picture is profoundly modified when one considers the new emerging paradigm of collective effects on supernova neutrino flavor conversions.

Recently, it was pointed out [23, 24, 25] that large neutrino density near the neutrinosphere results in significant coherent neutrino–neutrino forward scattering, which gives rise to collective neutrino flavor oscillations inside the SN. Three-flavor analysis of collective effects has now been carried out [26], which allows us to characterize collective oscillation effects and to write down the flavor spectra of neutrinos and antineutrinos arriving at the Earth. Following [26], we work in the modified flavor basis (ν_e, ν_x, ν_y), defined such that $(\nu_e, \nu_x, \nu_y) = R^\dagger_{23}(\theta_{23})(\nu_e, \nu_\mu, \nu_\tau)$, where R_{23} is the 2-3 rotation matrix. As a result of these collective oscillations, there are collective pair conversions $\nu_e\bar{\nu}_e \leftrightarrow \nu_y\bar{\nu}_y$ within the first $\mathcal{O}(100)$ km [24].

The manifestation of these flavor transformations depends on the primary spectra of neutrinos. In typical supernova models one finds a hierarchy of number fluxes $N_{\nu_e} > N_{\bar{\nu}_e} > N_{\nu_x} = N_{\bar{\nu}_x}$ [27, 28]. Even if it is not obvious that this hierarchy is maintained also at late times, in the following we will assume it as our benchmark. This scenario has been extensively studied analytically as well as numerically, and gives straightforward predictions for neutrino flavor conversions. In the normal hierarchy the spectra remain unaffected by collective oscillations. In the inverted hierarchy, for any non-zero value of θ_{13} such that the adiabatic solution in [29] is valid, the end of collective oscillations is marked by a complete exchange of the e and y flavor spectra for $\bar{\nu}$. The ν spectra also get swapped, however only above a characteristic energy, giving rise to a split in the spectrum [29, 30]. This solution is valid for extremely small θ_{13}, as long as bipolar oscillations develop sufficiently [31] and the evolution is adiabatic. We have checked that the adiabatic solution remains valid at values of θ_{13} that are as low as 10^{-10}, for typical SN neutrino density profiles. As a consequence, neutrino fluxes which are further processed by MSW matter effects are significantly different for the two hierarchies, even for extremely small θ_{13} values. This sensitivity presents a novel possibility to determine the mass hierarchy at small θ_{13}. We must remark that qualitatively different primary neutrino spectra and/or yet undiscovered flavor effects may yield different predictions for flavor conversion and the analysis will have to be repeated appropriately.

Here we concentrate on the $\bar{\nu}_e$ spectra observable through inverse beta decay reactions $\bar{\nu}_e + p \to n + e^+$ at water Cherenkov detectors. In inverted hierarchy, MSW matter effects in SN envelope are characterized in terms of the level-crossing probability P_H [15, 32] of antineutrinos, which is in general a function of the neutrino energy and θ_{13}. In the following, we consider two extreme limits, $P_H \simeq 0$ when $\sin^2\theta_{13} \gtrsim 10^{-3}$ ("large"), and $P_H \simeq 1$ when $\sin^2\theta_{13} \lesssim 10^{-5}$ ("small").

While propagating through the Earth, the $\bar{\nu}_e$ and $\bar{\nu}_x$ spectra partially mix. The neutrino fluxes F_ν at the Earth surface for normal hierarchy, as well as for inverted hierarchy with large θ_{13}, are given in terms of the the primary fluxes F^0_ν by

$$\begin{aligned} F_{\bar{e}} &= \cos^2\theta_{12} F^0_{\bar{e}} + \sin^2\theta_{12} F^0_{\bar{x}} \ , \\ F_{\bar{x}} &= \sin^2\theta_{12} F^0_{\bar{e}} + \cos^2\theta_{12} F^0_{\bar{x}} \ . \end{aligned} \tag{1}$$

For inverted hierarchy with small θ_{13}, we have

$$\begin{aligned} F_{\bar{e}} &= \cos^2\theta_{12} F^0_{\bar{y}} + \sin^2\theta_{12} F^0_{\bar{x}} \approx F^0_{\bar{x}} \ , \\ F_{\bar{x}} &= \sin^2\theta_{12} F^0_{\bar{y}} + \cos^2\theta_{12} F^0_{\bar{x}} \approx F^0_{\bar{x}} \ . \end{aligned} \tag{2}$$

Earth effect can be taken into account by just mapping $\cos^2\theta_{12} \to P(\bar{\nu}_1 \to \bar{\nu}_e)$ and $\sin^2\theta_{12} \to 1 - P(\bar{\nu}_1 \to \bar{\nu}_e)$, where $P(\bar{\nu}_1 \to \bar{\nu}_e)$ is the probability that a state entering the Earth as mass eigenstate $\bar{\nu}_1$ is detected as $\bar{\nu}_e$ at the detector.

From Eqs. (1) and (2), one expects to observe Earth matter effect in normal hierarchy independently of θ_{13}, while in inverted hierarchy it is expected only at large θ_{13}. For small θ_{13} and inverted hierarchy, the $\bar{\nu}_e$ spectrum arriving at the Earth is identical to the $\bar{\nu}_x$ spectrum arriving at the Earth, so any oscillation effect among them is unobservable. This implies that if next generation neutrino experiments bound θ_{13} to be small, from the (non)observation of Earth matter effect we could identify the neutrino mass hierarchy.

A strategy to observe Earth matter signatures in neutrino oscillations is to compare the signal at two detectors. The difference between the $\bar{\nu}_e$ flux $F^D_{\bar{e}}$ at a shadowed detector and the $\bar{\nu}_e$ flux $F_{\bar{e}}$ at a detector that is not shadowed by the Earth can be written as

$$\Delta F = F^D_{\bar{e}} - F_{\bar{e}} = f_{\rm reg}(F^0_{\bar{e}} - F^0_{\bar{x}}) \ , \tag{3}$$

for normal hierarchy as well as for inverted hierarchy with large θ_{13}. Here $f_{\rm reg} = P(\bar{\nu}_1 \to \bar{\nu}_e) - \cos^2\theta_{12}$ is the Earth regeneration factor. In inverted hierarchy for small θ_{13}, we get $\Delta F = 0$. If the $\bar{\nu}$ trajectories cross only the Earth mantle, characterized by an approximately constant density, $f_{\rm reg}$ is simply given by [16]

$$f_{\rm reg} = -\sin 2\tilde{\theta}_{12} \sin(2\tilde{\theta}_{12} - 2\theta_{12}) \sin^2\left(\frac{\Delta\tilde{m}^2_\odot L}{4E}\right) \ , \tag{4}$$

where $\tilde{\theta}_{12}$ is the effective value of the antineutrino mixing angle θ_{12} in matter, $\Delta\tilde{m}^2_\odot$ is the solar mass squared difference in matter, and L is the path length in Earth. In Earth matter, we have $\sin 2\tilde{\theta}_{12} > 0$ and $\sin(2\tilde{\theta}_{12} - 2\theta_{12}) < 0$, which tells us that $f_{\rm reg} \geq 0$.

The flavor dependent primary neutrino spectra $\varphi_\nu(E)$ can be conveniently parametrized as [27]

$$\varphi_\nu(E) = \frac{(\alpha+1)^{(\alpha+1)}}{\Gamma(\alpha+1)} \left(\frac{E}{\langle E\rangle_\nu}\right)^\alpha \frac{e^{-(\alpha+1)E/\langle E\rangle_\nu}}{\langle E\rangle_\nu} \ , \tag{5}$$

where Γ is the Euler Gamma function, $\langle E\rangle_\nu$ is the average energy for the different neutrino species, and α is the spectral pinching parameter. The neutrino flux at the neutrinosphere can then be estimated to be

$$F^0_\nu = \frac{L_\nu}{\langle E\rangle_\nu} \varphi_\nu(E) \ , \tag{6}$$

where L_ν is the luminosity in the ν flavor. All SN models robustly predict $\langle E_{\bar{e}}\rangle < \langle E_{\bar{x}}\rangle \approx \langle E_{\bar{y}}\rangle$, as well as $\alpha_{\bar{e}} \approx \alpha_{\bar{x}} \approx \alpha_{\bar{y}}$. This implies that the sign of $(F^0_{\bar{e}} - F^0_{\bar{x}})$ is positive at low energies (before the crossing of the $\bar{\nu}_e$ and $\bar{\nu}_x$ spectra) and negative at higher energies.

The net result is that when we compare the antineutrino fluxes between a shadowed and an unshadowed detector, we will have $\Delta F > 0$ at low energies and $\Delta F < 0$ at high energies in the case of normal mass hierarchy, or in inverted mass hierarchy with large θ_{13}. In inverted hierarchy with small θ_{13}, one expects a ΔF compatible with zero.

To illustrate the above, we consider a galactic supernova explosion at a distance of 10 kpc, with luminosities $L_{\bar{x}} = L_{\bar{y}} = 0.8 L_{\bar{e}}$ and total emitted energy $E_B = 3 \times 10^{53}$ erg. We also choose $\langle E\rangle_{\bar{e}} = 15$ MeV, $\langle E\rangle_{\bar{x}} = \langle E\rangle_{\bar{y}} = 18$ MeV, and $\alpha = 3$, inspired by the results of the Garching simulations [27, 33]. We analyze the detection of the above signal using two large water Cherenkov detectors A and B of fiducial mass 0.4 megaton each, as proposed for upcoming experiments [34, 35, 36]. We compare the number of events in detector A, where neutrinos arrive after traversing $L = 8000$ km in Earth mantle with an approximately constant density $\rho = 4.5$ g/cm^3, with another detector B for which the supernova is not shadowed by the Earth ($L = 0$). The reference values for features of the detectors, e.g., energy resolution and interaction cross sections, are the same as in [37]. We choose $\Delta m^2_\odot = 8 \times 10^{-5}$ eV^2 and $\sin^2\theta_{12} = 0.29$ as the oscillation parameters relevant for the Earth matter effect.



FIG. 1: Plot of the ratio R defined in Eq. 7, as a function of the observable positron energy for normal hierarchy (left panel) and inverted hierarchy (right panel), with $\sin^2\theta_{13} \lesssim 10^{-5}$. For $\sin^2\theta_{13} \gtrsim 10^{-3}$, the ratio R will be identical to the left panel for either hierarchy.

We define

$$R \equiv (N_A - N_B)/N_B \tag{7}$$

as the difference between the number of $\bar{\nu}_e$ events at the shadowed detector and the unshadowed detector, normalized to the number of events at the unshadowed detector. In Figure 1, we plot the ratio R as a function of the measured positron energy $E_{\rm pos}$ for $\bar{\nu}_e$ in normal hierarchy (left panel) and inverted hierarchy (right panel) for $\sin^2\theta_{13} \lesssim 10^{-5}$. The error bars show the statistical error in R. In the other extreme case of $\sin^2\theta_{13} \gtrsim 10^{-3}$, both the normal and inverted hierarchy would correspond to the left panel.

Let us consider the scenario where θ_{13} is known to be small. From the figure, in normal hierarchy the ratio R is positive for $E_{\rm pos} \lesssim 25$ MeV and negative at higher energy. The low energy spectrum is dominated by statistical error, but for $E_{\rm pos} \gtrsim 30$ MeV the depletion of the signal with respect to the unshadowed detector is clearly visible, with $|R| \gtrsim 5\%$. On the other hand, in inverted hierarchy we find $R = 0$. The difference in the predictions of two hierarchies is significant and should be observable. Primary spectra taken from Livermore simulations [28], which predict a larger difference between $\bar{\nu}_e$ and $\bar{\nu}_x$ average energies, would show a more pronounced Earth effect. We emphasize that our method is based on a model independent signature which does not rely on fitting or extracting any parameters.

The comparison of the neutrino signal in two detectors is also possible using only a single megaton class water Cherenkov detector together with the km^3 ice Cherenkov detector IceCube at the South Pole [38]. Even though IceCube cannot reconstruct the neutrino spectrum at SN energies, the ratio of luminosities at these two detectors can be determined rather accurately, which will show about 5% time variation if Earth effect is indeed present [18]. Moreover, if a large scintillator detector [34] is built, its superior energy resolution would allow the observation of the modulations induced by the earth effect in the spectrum, without the need to compare the signal with another unshadowed detector [19].

The swap of the $\bar{\nu}$ spectra due to collective effects does not depend on the exact neutrino density profile as long as the propagation is adiabatic [29], whose validity we have checked for typical SN profiles and θ_{13} as low as 10^{-10}. Decoherence effects are highly suppressed due to the ν_e–$\bar{\nu}_e$ flux asymmetry [39], and other multi-angle effects also do not affect the net antineutrino conversions substantially [25]. Moreover, with an extremely small θ_{13}, the detailed matter density profile near the H resonance is immaterial, and the effects of density fluctuations or turbulence may safely be ignored. Therefore, one can make the following statements: (i) Observation of Earth matter effects cannot be explained in inverted hierarchy (ii) Nonobservation of Earth matter effects cannot be explained in normal hierarchy (unless the primary fluxes are almost identical). Our proposed method is thus quite robust, and would be able to identify the mass hiererchy. It is not only competitive with the long baseline strategy proposed in [11], but also offers an independent astrophysical resolution to the hierarchy determination problem.

If θ_{13} is known to be large, the hierarchy can be determined through a number of other observables in the SN burst itself: signatures of SN shock-wave propagation in the $\bar{\nu}_e$ signal [37, 40, 41], the ν_e signal during the neutronization burst [42], or the direct, albeit extremely challenging, observation of the spectral split in ν_e spectrum [31] at a large liquid Argon detector [43]. In fact the hierarchy may even be identified at the long baseline experiments. However in such a scenario, the Earth matter effects act as an evidence for collective flavor conversions, thus giving us confidence about our understanding of the processes happening in the core of the star.

To conclude, determination of the leptonic mixing angle θ_{13} and the neutrino mass hierarchy represent two of the next frontiers of neutrino physics. In this Letter, we have proposed a new possibility for identifying the neutrino mass hierarchy that works for extremely small values of θ_{13}, far beyond the sensitivity of current and future terrestrial neutrino experiments. The sensitivity of supernova neutrino oscillations to the mass hierarchy, for incredibly small values of θ_{13}, is a consequence of the collective neutrino oscillations that take place near the supernova core. These effects interchange the initial $\bar{\nu}_e$ and $\bar{\nu}_y$ spectra in the inverted hierarchy, which are then further processed by MSW effects in the SN envelope. This spectral swap can be revealed by comparing the event rate at a shadowed detector with that at an unshadowed detector. If neutrino oscillation experiments fail to determine the mass hierarchy, then this proposed method could represent the last hope to resolve this issue, provided that large water Cherenkov detectors are available at the time of the next galactic SN explosion. This perspective should be considered when choosing optimal detector locations for upcoming large neutrino detectors [44].

The observation of the Earth matter effects in the inverted hierarchy for large θ_{13} also constitutes a smoking gun signature of collective oscillations in a SN, that arise from the as yet unprobed neutrino-neutrino interactions. It is fascinating to realize that intriguing effects like collective neutrino oscillations, occurring in the deep regions of an exploding supernova, produce observable signatures at Earth and enable us to probe neutrino properties. This confirms once again the extreme importance of supernovae as laboratories for fundamental neutrino physics.

We thank Georg Raffelt for valuable comments on the manuscript, and Andreu Esteban-Pretel, Sergio Pastor and Ricard Tomàs for important criticisms on its first version. A.M. also would like to thank Enrique Fernandez-Martinez for interesting discussions about mass hierarchy determination in future laboratory experiments. In Munich, this work was partly supported by the Deutsche Forschungsgemeinschaft (grant TR-27 "Neutrinos and Beyond"), by the Cluster of Excellence "Origin and Structure of the Universe" (Garching and Munich) and by the European Union (contract No. RII3-CT-2004-506222). In Mumbai, partial support by a Max Planck India Partnergroup grant is acknowledged. A.M. acknowledges support by the Alexander von Humboldt Foundation, and kind hospitality at the Tata Institute for Fundamental Research during the development of this work.

[1] A. Bandyopadhyay, S. Choubey, S. Goswami, S. T. Petcov and D. P. Roy, "Neutrino Oscillation Parameters After High Statistics KamLAND Results," arXiv:0804.4857 [hep-ph].

[2] M. C. Gonzalez-Garcia and M. Maltoni, "Phenomenology with Massive Neutrinos," Phys. Rept. **460**, 1 (2008) [arXiv:0704.1800 [hep-ph]].

[3] G. L. Fogli *et al.*, "Observables sensitive to absolute neutrino masses: A reappraisal after WMAP-3y and first MINOS results," Phys. Rev. D **75**, 053001 (2007) [hep-ph/0608060].

[4] M. Maltoni, T. Schwetz, M. A. Tortola and J. W. F. Valle, "Status of global fits to neutrino oscillations," New J. Phys. **6**, 122 (2004); The latest update is given in version 6 of hep-ph/0405172.

[5] A. Bandyopadhyay *et al.* [ISS Physics Working Group], "Physics at a future Neutrino Factory and super-beam facility," arXiv:0710.4947 [hep-ph].

[6] International Design Study of the Neutrino Factory web page, available at : www.hep.ph.ic.ac.uk/ids.

[7] R. Gandhi, P. Ghoshal, S. Goswami, P. Mehta, S. U. Sankar and S. Shalgar, "Mass Hierarchy Determination via future Atmospheric Neutrino Detectors," Phys. Rev. D **76**, 073012 (2007) [arXiv:0707.1723 [hep-ph]].

[8] T. Schwetz, "Determination of the neutrino mass hierarchy in the regime of small matter effect," JHEP **0705**, 093 (2007) [hep-ph/0703279].

[9] J. Learned, S. T. Dye, S. Pakvasa and R. C. Svoboda, "Determination of neutrino mass hierarchy and theta(13) with a remote detector of reactor antineutrinos," hep-ex/0612022.

[10] H. Minakata, H. Nunokawa, S. J. Parke and R. Zukanovich Funchal, "Determining neutrino mass hierarchy by precision measurements in electron and muon neutrino disappearance experiments," Phys. Rev. D **74**, 053008 (2006) [hep-ph/0607284].

[11] A. de Gouvea and W. Winter, "What would it take to determine the neutrino mass hierarchy if theta(13) were too small?," Phys. Rev. D **73**, 033003 (2006) [hep-ph/0509359].

[12] A. D. Bross, M. Ellis, S. Geer, O. Mena and S. Pascoli, "A Neutrino factory for both large and small θ_{13}," Phys. Rev. D **77**, 093012 (2008) [arXiv:0709.3889 [hep-ph]].

[13] L. Wolfenstein, "Neutrino oscillations in matter," Phys. Rev. D **17**, 2369 (1978).

[14] S. P. Mikheev and A. Y. Smirnov, "Resonance enhancement of oscillations in matter and solar neutrino spectroscopy," Sov. J. Nucl. Phys. **42**, 913 (1985) [Yad. Fiz. **42**, 1441 (1985)].

[15] A. S. Dighe and A. Y. Smirnov, "Identifying the neutrino mass spectrum from the neutrino burst from a supernova," Phys. Rev. D **62**, 033007 (2000) [hep-ph/9907423].

[16] C. Lunardini and A. Y. Smirnov, "Supernova neutrinos: Earth matter effects and neutrino mass spectrum," Nucl. Phys. B **616**, 307 (2001) [hep-ph/0106149].

[17] K. Takahashi, M. Watanabe and K. Sato, "The earth effects on the supernova neutrino spectra," Phys. Lett. B **510**, 189 (2001) [hep-ph/0012354].

[18] A. S. Dighe, M. T. Keil and G. G. Raffelt, "Detecting the neutrino mass hierarchy with a supernova at IceCube," JCAP **0306**, 005 (2003) [hep-ph/0303210].

[19] A. S. Dighe, M. T. Keil and G. G. Raffelt, "Identifying earth matter effects on supernova neutrinos at a single detector," JCAP **0306**, 006 (2003) [hep-ph/0304150].

[20] C. Lunardini and A. Y. Smirnov, "Probing the neutrino mass hierarchy and the 13-mixing with supernovae,"

JCAP **0306**, 009 (2003) [hep-ph/0302033].

[21] A. S. Dighe, M. Kachelriess, G. G. Raffelt and R. Tomas, "Signatures of supernova neutrino oscillations in the earth mantle and core," JCAP **0401** (2004) 004 [hep-ph/0311172].

[22] A. Dighe, "Neutrinos from a core collapse supernova," arXiv:0712.4386 [hep-ph].

[23] H. Duan, G. M. Fuller, J. Carlson and Y. Z. Qian, "Simulation of coherent non-linear neutrino flavor transformation in the supernova environment. I: Correlated neutrino trajectories," Phys. Rev. D **74**, 105014 (2006) [astro-ph/0606616].

[24] S. Hannestad, G. G. Raffelt, G. Sigl and Y. Y. Y. Wong, "Self-induced conversion in dense neutrino gases: Pendulum in flavour space," Phys. Rev. D **74** (2006) 105010 [Erratum-ibid. D **76** (2007) 029901] [astro-ph/0608695].

[25] G. L. Fogli, E. Lisi, A. Marrone and A. Mirizzi, "Collective neutrino flavor transitions in supernovae and the role of trajectory averaging," JCAP **0712**, 010 (2007) [arXiv:0707.1998 [hep-ph]].

[26] B. Dasgupta and A. Dighe, "Collective three-flavor oscillations of supernova neutrinos," Phys. Rev. D **77**, 113002 (2008) [arXiv:0712.3798 [hep-ph]].

[27] M. T. Keil, G. G. Raffelt and H. T. Janka, "Monte Carlo study of supernova neutrino spectra formation," Astrophys. J. **590**, 971 (2003) [astro-ph/0208035].

[28] T. Totani, K. Sato, H. E. Dalhed and J. R. Wilson, "Future detection of supernova neutrino burst and explosion mechanism," Astrophys. J. **496**, 216 (1998) [astro-ph/9710203].

[29] G. G. Raffelt and A. Y. Smirnov, "Self-induced spectral splits in supernova neutrino fluxes," Phys. Rev. D **76**, 081301 (2007) [Erratum-ibid. D **77**, 029903 (2008)] [arXiv:0705.1830 [hep-ph]].

[30] H. Duan, G. M. Fuller and Y. Z. Qian, "A Simple Picture for Neutrino Flavor Transformation in Supernovae," Phys. Rev. D **76**, 085013 (2007) [arXiv:0706.4293 [astro-ph]].

[31] H. Duan, G. M. Fuller, J. Carlson and Y. Q. Zhong, "Neutrino Mass Hierarchy and Stepwise Spectral Swapping of Supernova Neutrino Flavors," Phys. Rev. Lett. **99**, 241802 (2007) [arXiv:0707.0290 [astro-ph]].

[32] G. L. Fogli, E. Lisi, D. Montanino and A. Palazzo, "Supernova neutrino oscillations: A simple analytical approach," Phys. Rev. D **65**, 073008 (2002) [Erratum-ibid. D **66**, 039901 (2002)] [hep-ph/0111199].

[33] G. G. Raffelt, M. T. Keil, R. Buras, H. T. Janka and M. Rampp, 'Supernova neutrinos: Flavor-dependent fluxes and spectra," astro-ph/0303226.

[34] D. Autiero *et al.*, "Large underground, liquid based detectors for astro-particle physics in Europe: scientific case and prospects," JCAP **0711**, 011 (2007) [arXiv:0705.0116 [hep-ph]].

[35] C. K. Jung, "Feasibility of a next generation underground water Cherenkov detector: UNO," AIP Conf. Proc. **533**, 29 (2000) [hep-ex/0005046].

[36] K. Nakamura, "Hyper-Kamiokande: A next generation water Cherenkov detector," Int. J. Mod. Phys. A **18**, 4053 (2003).

[37] G. L. Fogli, E. Lisi, A. Mirizzi and D. Montanino, "Probing supernova shock waves and neutrino flavor transitions in next-generation water-Cherenkov detectors," JCAP **0504**, 002 (2005) [hep-ph/0412046].

[38] F. Halzen, "Astroparticle physics with high energy neutrinos: From AMANDA to IceCube," Eur. Phys. J. C **46**, 669 (2006) [astro-ph/0602132].

[39] A. Esteban-Pretel, S. Pastor, R. Tomas, G. G. Raffelt and G. Sigl, "Decoherence in supernova neutrino transformations suppressed by deleptonization," Phys. Rev. D **76**, 125018 (2007) [arXiv:0706.2498 [astro-ph]].

[40] R. Tomas, M. Kachelriess, G. Raffelt, A. Dighe, H. T. Janka and L. Scheck, "Neutrino signatures of supernova shock and reverse shock propagation," JCAP **0409**, 015 (2004) [astro-ph/0407132].

[41] G. L. Fogli, E. Lisi, A. Mirizzi and D. Montanino, "Damping of supernova neutrino transitions in stochastic shock-wave density profiles," JCAP **0606**, 012 (2006) [hep-ph/0603033].

[42] M. Kachelriess, R. Tomas, R. Buras, H. T. Janka, A. Marek and M. Rampp, "Exploiting the neutronization burst of a galactic supernova," Phys. Rev. D **71**, 063003 (2005) [astro-ph/0412082].

[43] I. Gil-Botella and A. Rubbia, "Oscillation effects on supernova neutrino rates and spectra and detection of the shock breakout in a liquid argon TPC," JCAP **0310**, 009 (2003) [hep-ph/0307244].

[44] A. Mirizzi, G. G. Raffelt and P. D. Serpico, "Earth matter effects in supernova neutrinos: Optimal detector locations," JCAP **0605**, 012 (2006) [astro-ph/0604300].